ASTHMA IN INDIANA



CHRONIC RESPIRATORY DISEASE SECTION

Champ Thomaskutty, MPH—*Director*, *Chronic Disease Epidemiology* Indiana State Department of Health

Overview

- □ Approximately 1 in 11 (9.1%) Indiana adults (18+ years) are currently diagnosed with asthma.
- □ Adult females in Indiana are more likely than males to have asthma.
- □ Approximately 1 in 10 (9.5%) Indiana children (ages 0-17) are currently diagnosed with asthma.
- Black children have a significantly higher prevalence rate than white children.
- □ An asthma diagnosis is more likely in adults with an annual household income < \$15,000 compared to household incomes ≥ \$15,000.




Most Common Data Sources

- □ Indiana Behavioral Risk Factor Surveillance System (BRFSS)
 - www.in.gov/isdh/22860.htm
- □ Centers for Disease Control and Prevention (CDC) BRFSS
 - www.cdc.gov/brfss
- □ BRFSS Asthma Call-Back Survey (ACBS)
 - http://www.cdc.gov/brfss/acbs/index.htm
- Indiana Hospital Discharge Database
 - http://www.state.in.us/isdh/20624.htm
- □ Indiana Vital Records
 - Information: <u>www.in.gov/isdh/23574.htm</u>
 - Data: www.in.gov/isdh/19096.htm



Prevalence of current asthma among adults (18+ years), Indiana, 2000–2010, 2011–2012*




*Starting in 2011 prevalence estimates were determined using a new, more precise methodology, including the addition of cell phone respondents and new weighting techniques; therefore, estimates made in 2011 or later should not be compared to earlier prevalence estimates.

Department of Health



Prevalence of current asthma among children (0-17 years), Indiana, 2005–2010, 2011*



*The 2011 prevalence estimate was determined using a new, more precise methodology, including the addition of cell phone respondents and new weighting techniques; therefore, the 2011 estimate should not be compared to earlier prevalence estimates.



Asthma self-management education among Hoosiers with current asthma

Has a health care professional	Adults ¹	Children ²
shown you how to use an inhaler?	93.7%	95.3%
taught you what to do during an asthma episode or attack?	78.4%	85.4%
watched you use an inhaler?	76.8%	82.7%
taught you how to recognize early signs or symptoms of an asthma episode?	67.1%	81.8%
taught you how to use a peak flow meter to adjust daily medications?	47.0%	39.7%
given you an asthma action plan?	35.5%	47.7%
Have you		
ever taken a course to manage asthma	8.8%	18.3%

^{1.} CDC and ISDH DAT. (2013). Behavioral Risk Factor Surveillance System Adult Asthma Call-back Survey, 2012.

^{2. &}lt;u>CDC</u> and <u>ISDH DAT</u>. (2012). Behavioral Risk Factor Surveillance System Child Multi-year Asthma Call-back Survey, 2006-2010.


Barriers facing Hoosiers with current asthma

	Adults ¹	Children ²
Uninsured	16.5%	4.1%
Unable to see primary care doctor for asthma because of the cost	15.9%	6.9%
Unable to see specialist because of the cost	9.8%	5.0%
Unable to buy medications because of the cost	30.6%	11.9%
Missed at least 1 day of work/school in the past 12 months due to asthma	33.6%	44.7%
Limited usual activities 'a lot' within the past 12 months due to asthma	9.6%	4.2%
Had no asthma symptom-free days during the past 2 weeks	29.9%	11.6%

^{1.} CDC and ISDH DAT. (2013). Behavioral Risk Factor Surveillance System Adult Asthma Call-back Survey, 2012.

^{2. &}lt;u>CDC</u> and <u>ISDH DAT</u>. (2012). Behavioral Risk Factor Surveillance System Child Multi-year Asthma Call-back Survey, 2006-2010.


Asthma control among adults (18+ years), Indiana, 2011






Source: <u>CDC</u> and <u>ISDH DAT</u>. (2013). *Behavioral Risk Factor Surveillance System Adult Asthma Call-back Survey*, 2012.

Asthma control among children (0–17 years), Indiana, 2006–2010





Asthma* ED visit rates,** Indiana, 2004–2012




^{*}Primary diagnosis asthma emergency department visits.

^{**}Age-adjusted rates per 10,000 population.





Asthma* ED visit rates** by month and age, Indiana, 2010–2012




^{*}Primary diagnosis asthma emergency department visits.



^{**}Age-specific rates per 10,000 population.

Asthma* hospitalization rates,** Indiana, 2004–2012




^{*}Primary diagnosis asthma hospitalizations.



^{**}Age-adjusted rates per 10,000 population.

Asthma* hospitalization rates** by month and age, Indiana, 2010–2012




^{*}Primary diagnosis asthma hospitalizations.




^{**}Age-specific rates per 10,000 population.

Emergency Department Visits, Overall and Children 5–17

Asthma Emergency Department Visits, Indiana, 2012




Child Asthma Emergency Department Visits, Indiana, Children 5–17 years, 2008–2012






In-patient Hospitalization, Overall and Children 5–17

Asthma Hospitalizations, Indiana, 2012



Child Asthma Hospitalizations, Indiana, Children 5–17 years, 2008–2012



For more information

- □ ISDH Asthma Program website
 - http://www.in.gov/isdh/24965.htm
- Most current Indiana data
 - http://www.in.gov/isdh/17279.htm
- □ Data request form
 - http://www.in.gov/isdh/files/Asthma ISDH Data Request Form.docx
- Indiana INdicators: health-related indicators for the state and by county
 - http://indianaindicators.org/



Contact Information

Champ Thomaskutty

Director, Chronic Disease Epidemiology Indiana State Department of Health cthomaskutty@isdh.in.gov (317) 233-7542

For more asthma data, visit the Asthma Program's <u>Data and Reports</u> webpage at http://www.in.gov/isdh/17279.htm.